

## Advances in Electron Donor Amendments

Short Course: Optimization and Monitoring for Remediation of Chlorinated and Related Compounds

Friday, November 22, 2019



## **About Us**




We research, develop, and commercialize innovative soil and groundwater remediation solutions in order to meet the increasing technological demands at contaminated sites.



## Soil and groundwater remediation of:

Chlorinated Solvents


- Petroleum Hydrocarbons
- NAPL Recovery



- 2 Saturated Zone NAPL Treatment
- 3 Dissolved Contaminant Anaerobic Remediation
- 4 Dissolved Contaminant Aerobic Remediation
- Implementation and Monitoring Systems




## **How Does Bioremediation Work?**





## **Biological Reductive Dechlorination**








### **Bioremediation Mechanisms**


#### Anaerobic Reductive Dechlorination






Modified from USGS WRI 99-2485

- Organic substrates that ferment to:
  - Acetate
  - Hydrogen (H<sub>2</sub>)
  - Hydrogen concentrations > 1 nM





- Organic substrates that ferment to:
  - Acetate
  - Hydrogen (H<sub>2</sub>)
  - Hydrogen concentrations > 1 nM
- Strongly reducing conditions
  - Sulfate Reducing or Methanogenic






- Organic substrates that ferment to:
  - Acetate
  - Hydrogen (H<sub>2</sub>)
  - Hydrogen concentrations > 1 nM
- Strongly reducing conditions
  - Sulfate Reducing or Methanogenic
- Right halorespiring bacteria
  - Dehalococcoides for DCE / VC



Dehalobacter restrictus



Dehalococcoides mccartyi Strain 195


Dhc = *Dehalococcoides* 

Dhb = *Dehalobacter* 

Other = Desulfitobacterium, sulfurospirillum, Clostridium



- Organic substrates that ferment to:
  - Acetate
  - Hydrogen (H<sub>2</sub>)
  - Hydrogen concentrations > 1 nM
- Strongly reducing conditions
  - Sulfate Reducing or Methanogenic
- Right halorespiring bacteria
  - Dehalococcoides for DCE / VC
- Nutrients
  - Vitamins and trace minerals to stimulate Dehalococcoides growth







### **Electron Donors**

 Average Composition and Electrons Released during Anaerobic Fermentation

	Atoms per Mole Substrate					Moles H2
Electron Donor	Carbon	Hydrogen	Oxygen	Average Molecular Weight	H2 Released per mole Substrate	released per gram substrate
Acetate	2	4	2	60.1	4	0.0666
Lactate	3	6	3	90.1	6	0.0666
Glucose	6	12	6	180.2	12	0.0666
Soybean Oil	56.3	99.5	6	873.1	156.5	0.1792



### Anaerobic Fermentation

 Soybean oil ferments to acetic acid and hydrogen

 $C_{56}H_{100}O_6$  (soybean oil<sup>1</sup>) + 50  $H_2O - B - >$ 

- $\underline{B}$ -> 28 CH<sub>3</sub>COOH (acetic acid) + 44 H<sub>2</sub>




<sup>1</sup>Represents weighted average of constituent fatty acids and glycerol.



## Soybean Fatty Acid Distribution

Fa	Percent	
C-16:0	Palmitic	11.0 %
C-18:0	Stearic	4.0 %
C-18:1	Oleic	24.0 %
C-18:2	Linoleic	54.0 %
C-18:3	Linolenic	7.0 %





## Why choose an EVO?

- Easily dispersed with groundwater
   (Oil-in-water emulsions are miscible with water)
- Low permeability loss
- Easy to implement
- Non-Toxic food-grade substance
- Limited chlorinated solvent sequestration




Oil in water emulsion, EDS-ER




### What is a surfactant?

- Molecule fits between oil and water
- Common and safe
- Found in
  - Salad dressing
  - Toothpaste
  - Mouthwash
  - Shampoo
- Must contact NAPL to work






## **Emulsifying Agents**



Vegetable Oil (Dispersed Phase)





## High Energy Shear Mixing










#### **Surfactant-Oil-Water Systems Phase Behavior**



# EDS-ERTM

# Electron Donor Solution – Extended Release

Water soluble vegetable oil









# "Greening" the cleanup

#### • EDS-ER:

- ✓ Eliminates Mechanical Energy inputs
- ✓ Allows Bulk Storage (long shelf life) and intermodal transportation
- ✓ Reduces need for excess drums and totes
- TASK™ EVO Self-Emulsifier
  - ✓ Easy Field Mixing
  - √ Source Local Soybean Oil
  - ✓ Reduced Carbon Footprint




## Field Mixing

• TASK™ EVO Self-Emulsifier


RBD Soybean Oil







## Impact of pH on Dechlorination



- pH of 6.0-8.5 is generally required for dechlorination to ethene\*
- pH 6.8-7.5 is considered optimal range, 7.5 is best\*
- Sites with low pH more likely to accumulate cDCE/VC



## Why is low pH so Common?

- Some sites have intrinsic groundwater pH in the 5.0-6.0 range
- Reductive dechlorination produces hydrochloric acid

 Fermentation of many electron donors produces acidic by products such as acetic acid


Lactic Acid 
$$\longrightarrow$$
 2H<sub>2</sub> + Acetate + CO<sub>2</sub>

+ CO<sub>2</sub> dissolves in water forming carbonic acid

Acid Generation
During
Bioremediation



# Ethene Production using KB-1® and Low pH KB-1® Plus at pH 6.0 and pH 7.0




Ethene production rate of low pH KB-1® Plus is 5 times higher than standard KB-1® at pH 6.0

Slide Courtesy of SiREM



## Project Approach



48 gal EVO



Products & Services for *In Situ* Remediation • tersusenv.com

## Field Application






EDS-ER™ Injections Using Water Powered Chemical Dosing Pump





Time Lapse Animation of the Dechlorination of a TCE Plume After Injections of *EDS-ER™* 





### Interested in a Site Evaluation?

#### www.tersusenv.com/support

#### **Immediate response:**

Sherri Scott

919.453.5577 x2003

919.527.9781 (mobile/text)

sherri.scott@tersusenv.com

- Options, Purpose & Due Date
- Tell Us About Your Site
  - Controlling Contaminant
  - Project Approach
  - Treatment Zone Physical Dimensions
  - Treatment Zone Hydrogeologic Properties
  - Aquifer Geochemistry
  - Natural Attenuation Parameters (not all applicable for each site)



## Sales and Technical Support



Tersus Environmental 919.453.5577 x2003 info@tersusenv.com

